Approved 28/04/2021

SCHEDULING STATUS S3

1 NAME OF THE MEDICINE
TELEASE 40 tablets
TELEASE 80 tablets

2 QUALITATIVE AND QUANTITATIVE COMPOSITION
TELEASE 40: Each tablet contains 40 mg telmisartan. Contains mannitol: 166.64 mg per tablet.
TELEASE 80: Each tablet contains 80 mg telmisartan. Contains mannitol: 333,28 mg per tablet. For the full list of excipients, see section 6.1.

3 PHARMACEUTICAL FORM
TELEASE 40: White or off-white to yellowish oblong biconvex uncoated tablets with break-line on one side and plain on the other side.
TELEASE 80: White or off-white to yellowish oval biconvex uncoated tablets with break-line on one side and plain on other side.

4 CLINICAL PARTICULARS
4.1 Therapeutic indications
Treatment of mild to moderate hypertension, either alone or in combination with hydrochlorothiazide.Reduction of cardiovascular morbidity and mortality in patients 55 years or older at high risk of cardiovascular disease; the benefit of treatment is evident after at least 6 months of continued treatment.
4.2 Posology and method of administration
Posology
Adults
Treatment of essential hypertension
The recommended dose is 40 mg once daily. In cases where the target blood pressure is not achieved, the TELEASE dose can be increased to a maximum of 80 mg once daily. Alternatively, TELEASE may be used in combination with a low dose thiazide diuretic such as hydrochlorothiazide 12,5 mg, which has been shown to have an additive blood pressure lowering effect with TELEASE. When considering raising the dose, it must be borne in mind that the maximum antihypertensive effect is generally attained four to eight weeks after the start of treatment.
Reduction of cardiovascular morbidity and mortality
The recommended dose is 80 mg once daily. It is not known whether doses lower than 80 mg of TELEASE are effective in preventing cardiovascular morbidity and mortality. When initiating TELEASE therapy for the prevention of cardiovascular morbidity and mortality, monitoring of blood pressure is recommended and, if appropriate, adjustment of medications that lower blood pressure may be necessary. The benefit of treatment is evident only after 6 months of continued treatment.
Renal impairment: No dosage adjustment is required for patients with renal impairment, including those on haemodialysis. TELEASE is not removed from blood by hemofiltration.
Hepatic impairment: In patients with mild to moderate hepatic impairment the dosage should not exceed 40 mg once daily.
Elderly: No dosage adjustment is necessary.
Children and adolescents up to 18 years: The safety and efficacy of TELEASE for use in children below 18 years have not been established.
Method of administration
TELEASE tablets are for once-daily oral administration and should be taken with liquid, with or without food.
4.3 Contraindications
- Hypersensitivity to telmisartan or to any of the ingredients of TELEASE (see section 2 and 6.1).
- A history of angioedema related to previous therapy with angiotensin-converting enzyme (ACE) inhibitors or angiotensin receptor blockers (ARBs). These patients must never again be given these medicines.
- Hereditary or idiopathic angioedema.
- Hypertrophic obstructive cardiomyopathy (HOCM).
- Severe renal function impairment (creatinine clearance less than 30 mL/min).
- Bilateral renal artery stenosis.
- Renal artery stenosis in patients with a single kidney.
- Aortic stenosis.
- Concomitant therapy with potassium sparing diuretics such as spironolactone, triamterene, amiloride (see section 4.4).
- Porphyria.
- Lithium therapy: Concomitant administration with TELEASE may lead to toxic blood concentrations of lithium (see section 4.5).
- Pregnancy and lactation (see section 4.6).
- Severe hepatic impairment.
- Obstructive biliary disorders.
- The concomitant use of TELEASE with aliskiren-containing products is contraindicated (see section 4.4 and 4.5).
- Concomitant use of fluoroquinolones with Angiotensin-converting enzyme (ACE) inhibitors/Angiotensin receptor blockers (ARBs) is contraindicated in patients with moderate to severe renal impairment (Creatinine Clearance ≤ 30 mL/min) and in elderly patients.

4.4 Special warnings and precautions for use
Should a woman become pregnant while receiving TELEASE, the treatment should be stopped promptly and switched to a different class of antihypertensive medicine (see sections 4.3 and 4.6).
Renovascular hypertension: There is an increased risk of severe hypotension and renal insufficiency when patients with bilateral renal artery stenosis or stenosis of the artery to a single functioning kidney are treated with TELEASE (see section 4.3).
Renal impairment and kidney transplant: When TELEASE is used in patients with impaired renal function, a periodic monitoring of potassium and creatinine serum levels is recommended. There is no experience regarding the administration of TELEASE in patients with a recent kidney transplant (see section 4.3).
Intravascular volume depletion: Symptomatic hypotension, especially after the first dose, may occur in patients who are volume and/or sodium depleted by vigorous diuretic therapy, dietary salt restriction, diarrhoea or vomiting. Such conditions, especially volume and/or sodium depletion, should be corrected before the administration of TELEASE.
Dual blockade of the renin-angiotensin-aldosterone system (RAAS): There is evidence that the concomitant use of angiotensin-converting enzyme (ACE) inhibitors, angiotensin II receptor blockers or aliskiren may increase the risk of hypotension, hyperkalaemia and decreases renal function (including acute renal failure). Dual blockade of RAAS through the combined use of TELEASE and aliskiren is therefore contraindicated (see section 4.3). TELEASE should not be used concomitantly with aliskiren (see section 4.3). ACE-inhibitors and angiotensin II receptor blockers should not be used concomitantly in patients with diabetic nephropathy.
Other conditions with stimulation of the renin-angiotensin-aldosterone system: In patients whose vascular tone and renal function depend predominantly on the activity of the renin-angiotensin-aldosterone system (e.g. patients with severe congestive heart failure or underlying renal disease, including renal artery stenosis), treatment with medicines that affect this system, such as TELEASE, has been associated with acute hypotension, uraemia, oliguria, or acute renal failure.
Concomitant use of fluoroquinolones: Concomitant use of fluoroquinolones and Angiotensin-converting enzyme (ACE) inhibitors/Angiotensin receptor blockers (ARBs) may precipitate acute kidney injury in patients, especially those with moderate to severe renal impairment and elderly patients (see section 4.3.) Renal function should be assessed before initiating treatment and monitored during treatment with fluoroquinolones or Angiotensin-converting enzyme (ACE) inhibitors/Angiotensin receptor blockers (ARBs) whether used separately and/or concomitantly.
Primary aldosteronism: Patients with primary aldosteronism generally will not respond to TELEASE. Therefore, the use of TELEASE is not recommended.
Aortic and mitral valve stenosis, obstructive hypertrophic cardiomyopathy: TELEASE is contraindicated in patients suffering from aortic or mitral stenosis, or obstructive hypertrophic cardiomyopathy (see section 4.3.). Excessive reduction in blood pressure in patients with ischaemic cardiopathy or ischaemic cardiovascular disease could result in a myocardial infarction or stroke.
Diabetic patients treated with insulin or antidiabetics: In these patients hypoglycaemia may occur with TELEASE treatment. Therefore, in these patients, appropriate blood glucose monitoring should be considered; a dose adjustment of insulin or antidiabetics may be required, when indicated.
Hyperkalaemia: During treatment with TELEASE, hyperkalaemia may occur, especially in the presence of renal impairment and/or heart failure. Monitoring of serum potassium in patients at risk is recommended. In the elderly, in patients with renal insufficiency, in diabetic patients, in patients concomitantly treated with other medicines that may increase potassium levels, and/or in patients with intercurrent events, hyperkalaemia may be fatal.
The main risk factors for hyperkalaemia are:
- Diabetes mellitus, renal impairment, age (>70 years)
- Combination with one or more other medicines that affect the renin-angiotensin-aldosterone system and/or potassium supplements. Medicines or therapeutic classes of medicines that may provoke hyperkalaemia are salt substitutes containing potassium, potassium-sparing diuretics, ACE inhibitors, angiotensin II receptor antagonists, non-steroidal anti-inflammatory medicines (NSAIDs, including selective COX-2 inhibitors), heparin, immunosuppressives (ciclosporin or tacrolimus), and trimethoprim.
- Intercurrent events, in particular dehydration, acute cardiac decompensation, metabolic acidosis, worsening of renal function, sudden worsening of the renal condition (e.g. infectious diseases), cellular lysis (e.g. acute limb ischemia, rhabdomyolysis, extend trauma).

Hepatic impairment: TELEASE is mostly eliminated in the bile. Patients with biliary obstruction disorders, cholestasis, or hepatic insufficiency can be expected to have reduced clearance (see section 4.3) and should not take TELEASE. TELEASE should be used only with caution in patients with mild to moderate hepatic impairment.
Other: Angiotensin receptor blockers, including TELEASE, are apparently less effective in lowering blood pressure in black people than in non-blacks, possibly because of higher prevalence of low-renin states in the black hypertensive population.

4.5 Interaction with other medicines and other forms of interaction
TELEASE may increase the hypotensive effect of other antihypertensive medicines. Co-administration of telmisartan did not result in a clinically significant interaction with warfarin, hydrochlorothiazide, glibenclamide, paracetamol, ibuprofen, simvastatin and amlodipine.
Digoxin: Co-administration of telmisartan did not result in a clinically significant interaction with digoxin, warfarin, hydrochlorothiazide, glibenclamide, paracetamol, ibuprofen, simvastatin and amlodipine. When telmisartan was co-administered with digoxin, median increases in digoxin peak plasma concentration (49 %) and in trough concentration (20 %) were observed (in a single dose study). When initiating, adjusting, and discontinuing telmisartan, monitor digoxin levels in order to maintain levels within the therapeutic range.
Potassium sparing diuretics or potassium supplements: As with other medicines acting on the renin-angiotensin-aldosterone system, telmisartan may provoke hyperkalaemia (see section 4.4). The risk may increase in case of treatment combination with other medicines that may also provoke hyperkalaemia (salt substitutes containing potassium, potassium-sparing diuretics, ACE inhibitors, angiotensin II receptor antagonists, non-steroidal anti-inflammatory medicines [NSAIDs, including selective COX-2 inhibitors], heparin, immunosuppressives [ciclosporin or tacrolimus], and trimethoprim).Angiotensin II receptor antagonists such as telmisartan, attenuate diuretic induced potassium loss. Potassium sparing diuretics e.g. spironolactone, eplerenone, triamterene, or amiloride, potassium supplements, or potassium-containing salt substitutes may lead to a significant increase in serum potassium.
Lithium: Reversible increases in serum lithium concentrations and toxicity have been reported during concomitant administration of lithium with angiotensin converting enzyme inhibitors. Increased serum levels have also been reported with telmisartan. Careful monitoring of serum lithium levels is recommended during concomitant use.
Non-steroidal anti-inflammatory drugs: Concomitant treatment with NSAIDs (including acetylsalicylic acid at anti- inflammatory dosage regimens, COX-2 inhibitors and non-selective NSAIDs) may reduce the antihypertensive effect of angiotensin II receptor antagonists. In some patients with compromised renal function (e.g. dehydrated patients or elderly patients with compromised renal function), the co-administration of angiotensin II receptor antagonists and medicines that inhibit cyclo-oxygenase may result in further deterioration of renal function, including possible acute renal failure, which is usually reversible. Therefore, the combination should be administered with caution, especially in the elderly. Compounds acting on the Renin-Angiotensin-System like TELEASE may have synergistic effects. Patients receiving NSAIDs and TELEASE should be adequately hydrated and be monitored for renal function at the beginning of combined treatment. A reduced effect of antihypertensive medicines like TELEASE by inhibition of vasodilating prostaglandins has been reported during combined treatment with NSAIDs.
Ramipril: In one study the co-administration of telmisartan and ramipril led to an increase of up to 2,5 fold in the AUC0-24 and Cmax of ramipril and ramiprilat. The clinical relevance of this observation is not known.
Dual blockade of the RAAS with ARBs, ACE inhibitors or aliskiren: Clinical trial data has shown that dual blockade of the renin-angiotensin-aldosterone-system (RAAS) through the concomitant use of angiotensin-converting enzyme (ACE) inhibitors, angiotensin II receptor blockers or aliskiren is associated with a higher frequency of adverse events such as hypotension, hyperkalaemia and decreased renal function (see sections 4.3 and 4.4). Based on their pharmacological properties it can be expected that the following medicines may potentiate the hypotensive effects of all antihypertensives including telmisartan: Baclofen, amifostine; Antipsychotic hypotensive medicines; Tricyclic antidepressants, barbiturates, narcotics, or antidepressants.
Concomitant use of fluoroquinolones: Concomitant use of fluoroquinolones and Angiotensin-converting enzyme (ACE) inhibitors/Angiotensin receptor blockers (ARBs) may precipitate acute kidney injury. The mechanism of the possible interaction between the different classes of medicines, over and above different mechanisms of kidney damage, is unknown (see section 4.3).
Diuretics (thiazide or loop diuretics): Prior treatment with high dose diuretics such as furosemide (loop diuretic) and hydrochlorothiazide (thiazide diuretic) may result in volume depletion, and in a risk of hypotension when initiating therapy with telmisartan. To be taken into account with concomitant use.
Corticosteroids (systemic route): Reduction of the antihypertensive effect.
4.6 Fertility, pregnancy and lactation
Women of childbearing potential: Women of childbearing age should ensure effective contraception.
Pregnancy: Safety in pregnancy has not been established (see section 4.3). When pregnancy is planned or confirmed TELEASE should be discontinued. Medicines affecting the renin-angiotensin system, such as TELEASE, can cause embryonal toxicity, foetal and neonatal morbidity and mortality when administered to pregnant women.
Breastfeeding: TELEASE is contraindicated during lactation (see section 4.3).
Fertility: In preclinical studies, no effects of TELEASE on male and female fertility were observed
4.7 Effects on ability to drive and use machines
No studies on the effect on the ability to drive and use machines have been performed. However, when driving vehicles or operating machinery it should be taken into account that dizziness or drowsiness may occasionally occur when taking antihypertensive therapy including TELEASE.
4.8 Undesirable effects
Tabulated list of adverse reactions

MedDRA system organ class	Frequency	Adverse reactions
Infections and infestations	Less frequent	Urinary tract infection (including cystitis), upper respiratory tract infection including pharyngitis and sinusitis
	Frequency unknown	Sepsis including fatal outcome
Blood and lymphatic system disorders	Less frequent	Anaemia, thrombocytopenia
	Frequency unknown	Eosinophilia
Immune system disorders	Less frequent	Hypersensitivity, angio-oedema (with fatal outcome)
	Frequency unknown	Anaphylactic reaction
Metabolism and nutrition disorders	Less frequent	Hyperkalaemia
	Frequency unknown	Hypoglycaemia (in diabetic patients)
Psychiatric disorders	Less frequent	Insomnia, depression, anxiety
Nervous system disorders	Less frequent	Syncope/fainting, somnolence
Eye disorders	Less frequent	Visual disturbance
Ear and labyrinth disorders	Less frequent	Vertigo
Cardiac disorders	Less frequent	Bradycardia, tachycardia
Vascular disorders	Less frequent	Hypotension, orthostatic hypotension
Respiratory, thoracic and mediastinal disorders	Frequent	Cough
	Less frequent	Dyspnoea, interstitial lung disease
Gastrointestinal disorders	Less frequent	Abdominal pain, diarrhoea, dyspepsia, flatulence, vomiting, dry mouth, stomach discomfort, dysgeusia
Hepato-biliary disorders	Frequency unknown	Hepatic function abnormal/liver disorder
Skin and subcutaneous tissue disorders	Less frequent	Increased sweating (hyperhidrosis), pruritis, rash, eczema, erythema, drug eruption, toxic skin eruption
	Frequency unknown	Urticaria
Musculoskeletal and connective tissue disorders	Less frequent	Back pain (e.g. sciatica), muscle spasms (cramps in legs), myalgia, arthralgia, pain in extremity (leg pain)
	Frequency unknown	Tendon pain (tendinitis like symptoms)
Renal and urinary disorders	Less frequent	Renal impairment including acute renal failure
General disorders and administration site conditions	Less frequent	Chest pain, asthenia (weakness), influenza-like symptoms
Investigations	Less frequent	Blood creatinine increased, haemoglobin decreased, blood uric acid increased, hepatic enzymes increased, blood creatine phosphokinase increased

Reporting of suspected adverse reactions: Reporting suspected adverse reactions after authorisation of the medicine is important. It allows continued monitoring of the benefit/risk balance of the medicine. Health care providers are asked to report any suspected adverse reactions to SAHPRA via the "**6.04 Adverse Drug Reactions & Quality Problem Reporting Form**", found online under SAHPRA's publications: https://sahpra.org.za/wp-content/uploads/2020/01/6.04_ARF1_v5.1_27Jan2020.pdf
4.9 Overdose
Symptoms: Limited information is available with regard to overdose in humans. The most prominent manifestations of telmisartan overdose were hypotension and tachycardia; bradycardia dizziness, increase in serum creatinine, and acute renal failure have also been reported.
Management: If symptomatic hypotension should occur, supportive treatment should be instituted. Telmisartan, as in TELEASE is not removed by haemodialysis. Management depends on the time since ingestion and the severity of the symptoms. Serum electrolytes and creatinine should be monitored frequently. If hypotension occurs, the patient should be placed in a supine position, with salt and volume replacement given quickly.

5 PHARMACOLOGICAL PROPERTIES
5.1 Pharmacodynamic properties
A.T.1.3 Vascular medicines – other hypotensives. Pharmacotherapeutic group: Angiotensin II Antagonists, plain, ATC Code: C09CA07.
Telmisartan is a specific angiotensin II receptor (type AT$_1$) antagonist. It displaces angiotensin II from its binding site at the AT$_1$ receptor subtype, which is responsible for the known actions of angiotensin II. Telmisartan does not exhibit any partial agonist activity at the AT$_1$ receptor. Telmisartan selectively binds at the AT$_1$ receptor. The binding is long-lasting. Telmisartan does not inhibit human plasma renin or block ion channels. In human, an 80 mg dose of telmisartan almost completely inhibits the angiotensin II evoked blood pressure increase. The inhibitory effect is maintained over 24 hours and is still measurable up to 48 hours. After administration of the first dose of telmisartan in hypertensive patients, onset of antihypertensive activity occurs within 3 hours. The maximum reduction in blood pressure is generally attained 4 weeks after the start of treatment and is sustained during long-term therapy. The antihypertensive effect persists over 24 hours after dosing. There is an apparent trend to a dose relationship with regard to a time to recovery of baseline systolic blood pressure. In this respect data concerning diastolic blood pressure are inconsistent. In patients with hypertension, telmisartan reduces both systolic and diastolic blood pressure without affecting pulse rate. Upon abrupt cessation of treatment with telmisartan, blood pressure gradually returns to pre-treatment values over a period of several days without evidence of rebound hypertension. Telmisartan treatment has been shown in clinical trials to be associated with statistically significant reductions in proteinuria (including microalbuminuria and macroalbuminuria) in patients with hypertension and diabetic nephropathy.Telmisartan treatment has been shown to be associated with statistically significant reductions in Left Ventricular Mass and Left Ventricular Mass Index in patients with hypertension and Left Ventricular Hypertrophy.
5.2 Pharmacokinetic properties
Absorption of telmisartan is rapid although the amount absorbed varies. The mean absolute bioavailability for telmisartan is about 50 %. When TELEASE is taken with food, the reduction in the area under the plasma concentration-time curve (AUC$_{0-\infty}$) of telmisartan varies from approximately 6 % (40 mg dose) to approximately 19 % (160 mg dose). After 3 hours post administration, plasma concentrations are similar whether telmisartan is taken fasting or with food. The small reduction in AUC is not expected to cause a reduction in the therapeutic efficacy. Gender differences in plasma concentrations were observed, C$_{max}$ and AUC being approximately 3- and 2-fold higher, respectively, in females compared to males without relevant influence on efficacy. Telmisartan is highly bound to plasma protein (> 99,5 %), mainly albumin and alpha-1 acid glycoprotein. The mean steady state apparent volume of distribution (V$_{ss}$) is approximately 500 L. Telmisartan is metabolised by conjugation to the glucuronide. No pharmacological activity has been shown for the conjugate. Telmisartan is characterised by biexponential decay pharmacokinetics with a terminal elimination half-life of > 20 hours. The maximum plasma concentration (C$_{max}$) and, to a smaller extent, area under the plasma concentration-time curve (AUC) increase disproportionately with dose. There is no evidence of clinically relevant accumulation of telmisartan. Plasma concentrations were higher in females than in males, without relevant influence on efficacy. After oral administration telmisartan is nearly exclusively excreted with the faeces, mainly as unchanged compound. Cumulative urinary excretion is < 2 % of dose. Total plasma clearance (CL$_{tot}$) is high (approximately 900 mL/min) when compared with hepatic blood flow (about 1 500 mL/min).
Special populations
Elderly patients: The pharmacokinetics of telmisartan do not differ between younger and elderly patients.
Patients with renal impairment: Lower plasma concentrations were observed in patients with renal insufficiency undergoing dialysis. Telmisartan is highly bound to plasma protein in renal-insufficient subjects and cannot be removed by dialysis. The elimination half-life is not changed in patients with renal impairment.
Patients with hepatic impairment: Pharmacokinetic studies in patients with hepatic impairment showed an increase in absolute bioavailability up to nearly 100 %. The elimination half-life is not changed in patients with hepatic impairment.

6 PHARMACEUTICAL PARTICULARS
6.1 List of excipients
Mannitol (E421), Meglumine, Povidone, Sodium hydroxide (E524), Sodium stearyl fumarate.
6.2 Incompatibilities
Not applicable.
6.3 Shelf life
36 months.
6.4 Special precautions for storage
Store at or below 25 °C. Keep the blisters in the original container until required for use.
6.5 Nature and contents of container
TELEASE tablets are packed in the following pack: Blister pack using aluminium foil with heat seal lacquer coating sealed with cold forming blister aluminium foil (Alu/Alu blister). The blister strips will be packed in cartons with 14, 28, 30 or 56 tablets. Not all pack sizes may be marketed.
6.6 Special precautions for disposal and other handling
Not applicable

7 HOLDER OF CERTIFICATE OF REGISTRATION
Trinity Pharma (Pty) Ltd. 106 16th Road, Midrand, 1686.

8 REGISTRATION NUMBERS
TELEASE 40: 47/7.1.3/0334
TELEASE 80: 47/7.1.3/0335

9 DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION
16 February 2021

10 DATE OF REVISION OF THE TEXT

SKEDULERINGSTATUS S3

1 HANDELSNAAM VAN DIE MEDISYNE
TELEASE 40 tablette
TELEASE 80 tablette

2 KWALITATIEWE EN KWANTITATIEWE SAMESTELLING
TELEASE 40: Elke tablet bevat 40 mg telmisartan. Bevat mannitol: 166,64 mg per tablet.
TELEASE 80: Elke tablet bevat 80 mg telmisartan. Bevat mannitol: 333,28 mg per tablet. Vir 'n volledige lys van hulpstowwe, sien afdeling 6.1.

3 DOSEERVORM
TELEASE 40: Wit of effektief wit geel, langwerpige, bikonvekse onbedekte tablet met 'n breuklyn gebosseleer aan die een kant en aan die ander kant 'n plein afwerking.
TELEASE 80: Wit of effektief wit geel, ovaalvormige, bikonvekse onbedekte tablet met 'n breuklyn gebosseleer aan die een kant en aan die ander kant 'n plein afwerking.

4 KLINIESE INLIGTING
4.1 Terapeutiese indikasies
Behandeling van ligte tot matige hipertensie, as enkelmiddel of in kombinasie met hidrochlortiasied. Verlaging van kardiovaskulêre morbiditeit en mortaliteit in pasiënte 55 jaar en ouer met 'n hoë risiko vir kardiovaskulêre siektes; die voordeel van die behandeling is duidelik na 6 maande van ononderbroke behandeling.

4.2 Posologie en metode van toediening
Posologie
Volwassenes
Behandeling van essensiële hipertensie: Die aanbevole dosis is 40 mg daagliks. In gevalle waar die gewenste bloeddruk nie verkry word nie, kan die TELEASE dosis verhoog word tot 80 mg daagliks. Alternatiewelik kan TELEASE in kombinasie met 'n lae dosis tiasied-tipe diuretikum soos hidrochlortiasied 12,5 mg, wat 'n aanvullende bloeddrukverlagende effek met TELEASE getoon het, toegedien word. Wanneer daar oorweeg word om die dosis te verhoog, moet daar in gedagte gehou word dat die maksimum antihipertensiewe effek oor die algemeen vier tot agt weke, na aanvang van behandeling, verkry word.
Vermindering van kardiovaskulêre morbiditeit en mortaliteit: Die aanbevole dosis is 80 mg daagliks. Dit is nie bekend of dosisse laer as 80 mg TELEASE doeltreffend is om kardiovaskulêre morbiditeit en mortaliteit te voorkom nie. Tydens die aanvang van behandeling met TELEASE vir die voorkoming van kardiovaskulêre morbiditeit en mortaliteit, word dit aanbeveel dat die bloeddruk van die pasiënt gemonitor word, en indien van toepassing, kan die mediseyne wat die bloeddruk verlaag sodoende aangepas word. Die voordeel van die behandeling is eers duidelik na 6 maande van ononderbroke behandeling.
Ingeskorte nierfunksie: Geen aanpassing in die dosis is nodig by pasiënte met ingeskorte nierfunksie nie, insluitend diegene wat op hemodialise is. TELEASE word nie deur hemofiltrasie uit die bloed verwyder nie.
Ingeskorte leverfunksie: Pasiënte met ligte tot matige leverontoereikendheid moet nie die dosis van 40 mg daagliks oorskry nie.
Bejaardes: Geen aanpassing in die dosis is nodig nie.
Kinders en adolessente tot en met 18 jaar: Die veiligheid en doeltreffendheid van TELEASE vir gebruik by kinders onder 18 jaar is nie nog nie vasgestel nie.
Metode van toediening: TELEASE tablette is vir een keer per dag orale toediening bestem en moet saam met vloeistof, met of sonder voedsel geneem word.
4.3 Kontraindikasies
- Hipersensitiwiteit vir telmisartan of enige van die ander bestandele in TELEASE (sien afdeling 2 en 6.1).
- 'n Geskiedenis van angioneurotiese edeem verwant aan voorafgaande behandeling met angiotensien-converting enzyme (ACE) inhibeerders of angiotensien-reseptor blokkeerders (ARBs): Daar moet nooit weer van die mediseyne vir hierdie pasiënte gegee word nie.
- Overerflike of idiopatiese angioedeem.
- Hipertrofiese obstruktiewe kardiomiopatie (HOKM).
- Ernstige belemmerde nierfunksie (kreatinienopruiming minder as 30 mL/min).
- Bilaterale nierarasstenose.
- Nierarasstenose by pasiënte met 'n enkele nier.
- Stenose van die aorta.
- Gelyktydige behandeling met kaliumsparende diuretika soos spironolaktoon, triamtereen, amiloried (verwys na afdeling 4.4).
- Porfirie.
- Litiumbehandeling: Gelyktydige toediening van TELEASE kan lei tot 'n toksiese bloedkonsentrasie van litium (verwys na afdeling 4.5).
- Swangerskap en laktasie (verwys na afdeling 4.4 en 4.6).
- Ernstige ingekorte lewerfunksie.
- Obstruksie galbuisafwykings.
- Die gelyktydige gebruik van TELEASE met aliskiren-bevattende produkte word teenaangedui (verwys na afdeling 4.4 en 4.5).
- Gelyktydige gebruik van fluorokinolone met Angiotensien-omskakelingsensiem (ACE) inhibeerders/Angiotensien reseptor blokkeerders (ARB's) is kontraindikasie in pasiënte met matige tot ernstige nierversaking (kreatinienopklaring van ≤ 30 mL/min) en in bejaarde pasiënte.

4.4 Spesiale waarskuwings en voorsorgmaatreëls vir gebruik
Indien 'n vrou swanger raak tydens TELEASE, moet die behandeling onmiddellik gestaak word en na 'n ander klas van anti-hipertensiewe middel oorgeskakel word (verwys na afdeling 4.3 en 4.6).
Renovaskulêre hipertensie: Daar is 'n hoër risiko van ernstige hipotensie en nierontoereikendheid wanneer pasiënte met bilaterale nierarteriestenose of stenose van die slagaar na 'n enkele funksionerende nier met TELEASE behandel word (verwys na afdeling 4.3).
Belemmerde nierfunksie en nieroorplanting: Wanneer TELEASE gebruik word in pasiënte met belemmerde nierfunksie, word periodieke monitering van die kalium- en kreatinienserumvlakke aanbeveel. Daar is geen ervaring met die toediening van TELEASE in pasiënte met 'n onlangse nieroorplanting nie (verwys na afdeling 4.3).
Intravaskulêre hipovolemie: Simptomatiese hipotensie, veral na die eerste dosis, kan voorkom by pasiënte met volume- en/of natriumuitputting weens drastiese diuretiese terapie, soutbeperkende dieet, diaree of braking. Suike toestande, veral volume- en/of natriumuitputting, moet voor die toediening van TELEASE reggestel word.
Dubbele blokkade van die renien-angiotensien-aldosteronsisteel (RAAS): Daar is bewyse dat die gelyktydige gebruik van angiotensien omskakelingsensiem (ACE) inhibeerders, angiotensien-II reseptorblokkeerders of aliskiren die risiko mag verhoog van hipotensie, hiperkalemie en verswakking van die nierfunksie (insluitend akute nierversaking). Dubbele blokkade van RAAS deur die gelyktydige gebruik van TELEASE en aliskiren word dus teenaangedui. TELEASE moet nie gelyktydig met aliskiren gebruik word nie (verwys na afdeling 4.3). ACE-inhibeerders en angiotensien-II reseptorblokkeerders moet nie gelyktydig gebruik word in pasiënte met diabetiese nefropatie nie.
Ander siektetoestande met stimulasie van die renien-angiotensien-aldosteronsisteel: By pasiënte wie se vaskulêre tonus en nierfunksie hoofsaaklik afhanklik is van die aktiwiteit van die renien-angiotensien-aldosteronsisteel (bv. pasiënte met ernstige kongestiewe hartversaking of onderliggende niersiekte, insluitend nierarasstenose), word die behandeling met geneesmiddels wat hierdie stelsel affekteer, soos TELEASE, geassosieer met akute hipotensie, oliguria of akute nierversaking.
Gelyktydige gebruik van fluorokinolone: Gelyktydige gebruik van fluorokinolone en Angiotensien-omskakelingsensiem (ACE) inhibeerders/ Angiotensien reseptor blokkeerders (ARBs) kan akute nierbesering in pasiënte bespoedig, veral die met matige tot ernstige nierversaking en bejaarde pasiënte (verwys na afdeling 4.3.) Nierfunksie moet bepaal word voor die instelling van behandeling en moet gemonitor word tydens behandeling met fluorokinolone of ACE inhibeerders/
Angiotensien reseptor blokkeerders (ARBs) hetsy afsonderlik of gesamentlik geneem word.
Primêre aldosteronisme: Pasiënte met primêre aldosteronisme sal oor die algemeen nie op TELEASE reageer nie. Gevolglik word die gebruik van TELEASE nie aanbeveel nie.
Aorta- en mitraalklep stenose, obstruktiewe hipertrofiese kardiomiopatie: TELEASE is teenaangedui in pasiënte wat ly aan aortiese- of mitralisstenose of hipertrofiese obstruktiewe kardiomiopatie (verwys na afdeling 4.3). Oormatige verlaging in bloeddruk in pasiënte met iskemiese kardiopatie of iskemiese kardiovaskulêre siekte mag ontaard in 'n miokardiale infarksie of beroerte.
Diabetiese pasiënte wat behandel word met insulien of antidiabetiese medisyne: Hipoglisemie mag voorkom by hierdie pasiënte **tydens behandeling met TELEASE**. Daarom moet daar gepaste bloedglukose monitering oorweeg word by hierdie pasiënte; 'n dosisaanpassing van insulin of antidiabetiese medisyne mag nodig wees, soos aangedui.
Hiperkalemie: Tydens die behandeling met TELEASE mag hiperkalemie voorkom, veral in die teenwoordigheid van belemmerde nierfunksie en/of hartversaking. Monitering van serumkalium
Lewerensiemversteuring: Telmisartan word hoofsaaklik in die galpyp deur middel van geabsorbeer word wissel. Die gemiddelde absolute biobeskikbaarheid is ongeveer 50 % vir telmisartan. Wanneer TELEASE met voedsel geneem word, wissel die vermindering in oppervlakte onder die plasmakonsentrasie-tydkromme (AUC$_{0-\infty}$) van telmisartan vanaf ongeveer 6 % (40 mg dosis) tot ongeveer 19 % (160 mg dosis). Drie (3) ure na toediening is die plasmakonsentrasies vergelykbaar, ongeag daarvan of telmisartan vastend of met voedsel geneem is. Daar word nie verwag dat die klein vermindering in AOK verminderde terapeutiese doeltreffendheid sal veroorsaak nie. Verskille in plasmakonsentrasies tussen geslagte word waargeneem; die K$_{max}$ en AOK is onderskeidelik ongeveer 3- en 2- keer hoër by vroue in vergelyking met mans, sonder 'n relevante invloed op doeltreffendheid. Telmisartan word sterk (> 99,5 %) aan plasmaproteïen gebind, hoofsaaklik albumien en alfa 1-suur-glikoproteïen. Die gemiddelde skynbare verspreidingsvolume (Vdss) by stabiele staat is ongeveer 500 L. Telmisartan word deur middel van konjugasie na die glukuronied gemetaboliseer. Die konjugaat toon geen farmakologiese aktiwiteit nie. Telmisartan word gekenmerk deur bi-eksponensiële afbraak farmakokinetika met 'n terminale eliminasiehalfleeftyd van > 20 uur. Die maksimum plasmakonsentrasie (K$_{max}$) en tot 'n mindere mate die area onder die plasma konsentrasietydskromme word nie-lineêr met die dosis. Daar is geen bewys van klinies relevante akkumulasie van telmisartan nie. Plasmakonsentrasies was hoër by vroue as by mans, sonder enige relevante invloed op die doeltreffendheid. Na mondelikse toediening word telmisartan feitlik net deur die feses uitgeskei, hoofsaaklik as die onveranderde produk. Die kumulatiewe hoeveelheid wat via die urine uitgeskei word is < 2 % van die dosis. Algehele opruiming uit die plasma (CL$_{tot}$) is hoog (ongeveer 900 mL/min) indien dit met die hepatiese bloedvloei (ongeveer 1 500mL/min) vergelyk word.
Spesiale pasiëntgroepe
Bejaarde pasiënte: Die farmakokinetika van telmisartan verskil nie tussen jonger en ouer pasiënte nie.
Pasiënte met ingekorte nierfunksie: Laer plasmakonsentrasies is by pasiënte met nierontoereikendheid wat dialise ondergaan waargeneem. By pasiënte met nierontoereikendheid is telmisartan baie sterk aan plasmaproteïene gebind en kan dit nie deur dialise verwyder word nie. Die eliminasiehalfleeftyd verander nie by pasiënte met belemmerde nierfunksie nie.
Pasiënte met belemmerde lewerfunksie: By pasiënte met lewerontoereikendheid is daar met farmakokinetiese studies 'n toename van bykans 100 % in die absolute biobeskikbaarheid waargeneem. Die eliminasiehalfleeftyd verander nie by pasiënte met lewerontoereikendheid nie.

6 FARMASEUTIESE BESONDERHEDE/INLIGTING
6.1 Lys van bymiddels
Mannitol (E421), Meglumien, Povidoon, Natriumhidroksied (E524) , Natriumstearielfumaraat.
6.2 Onverenigbaarheid
Nie van toepassing.
6.3 Rakleeftyd
36 maande.
6.4 Spesiale voorsorgmaatreëls tydens berging
Stoor teen of benede 25 °C. Hou die stulpstrook in die oorspronklike verpakking, tot voor gebruik.
6.5 Aard en inhoud van die houer
TELEASE tablette word verpak in die volgende verpakking: Stulpstroke van gewone silwer aluminiumfolie met 'n hitte verseelde laag bedek met 'n koue vormende aluminiumfolie stulpstrook. Die stulpstroke word verpak in kartonhouers van 14, 28, 30 or 56 tablette. Nie alle verpakkingsgroottes word bemark nie
6.6 Spesiale voorsorgmaatreëls vir die vegdoening en ander hantering
Nie van toepassing nie.

7 HOUER VAN DIE REGISTRASIESERTIFIKAAT
Trinity Pharma (Pty) Ltd. 16e Straat 106, Midrand, 1686

8 REGISTRASIENOMMERS
TELEASE 40: 47/7.1.3/0334
TELEASE 80: 47/7.1.3/0335

9 DATUM VAN DIE EERSTE GOEDKEURING/HERSIENING VAN DIE GOEDKEURING
16 Februarie 2021

10 DATUM VAN DIE HERSIENING VAN DIE TEKS

PATIENT INFORMATION LEAFLET

SCHEDULING STATUS S3

TELEASE 40 tablets
TELEASE 80 tablets

Telmisartan
Contains sugar: mannitol 166,64 mg per 40 mg tablet
Contains sugar: mannitol 333,28 mg per 80 mg tablet

Read all of this leaflet carefully before you take TELEASE
- Keep this leaflet. You may need to read it again.
- If you have further questions, please ask your doctor, pharmacist, nurse or other health care provider.
- TELEASE has been prescribed for you personally and you should not share your medicine with other people. It may harm them, even if their symptoms are the same as yours.

1. What is in this leaflet

1. What TELEASE is and what it is used for
2. What you need to know before you take TELEASE
3. How to take TELEASE
4. Possible side effects
5. How to store TELEASE
6. Contents of the pack and other information

**2. What TELEASE is and